PHYSICIAN'S ORDERS

Authorization is given to the Pharmacy Department to dispense a generic & therapeutic equivalent drug when a brand name is ordered unless checked here.

ALLERGIES:

7	.OILO.				
USE B	ALL POI	NT PEN - PLEASE BEAR DOWN			
	□ HC □ DN	ne HCG Result: FG QL Result: IA (1 purple for kit) xicology (2 gray for kit)	☐ Anti HB sAG	for kit)	
	√ BY RN	CHECK APPROPRIATE BOXES	S A N E - MD ORDER FORM		
		□ Antiemetic 30 minutes prior to Pregnancy/STD prophylaxis: Prochlorperazine (Compazine) 10 mg PO X1			
		PREGNANCY PROPHYLAXIS (within 72 hours of assault) Plan B (Levonorgestrel 0.75 mg) Dose: Take one (1) tablet as soon as possible within 72 hrs after unprotected intercourse. Take the second dose of one (1) tablet 12 hrs later.			
		STD PROPHYLAXIS			
		DISEASE	RECOMMENDED	ALTERNATIVE	
		GONORRHEA	 Ceftriaxone 250 mg IM Contraindication: Allergy to Penicillin/Cephalosporins 	☐ Ciprofloxacin 500 mg PO X1	
		CHLAMYDIA	 Azithromycin 1 gm PO X 1 Contraindication: Allergy to Erythromycin, Azithromycin or other macrolide antibiotics. 	☐ Doxycycline 100 mg PO BID X 7 days. -or- ☐ Erythromycin 500 mg PO QID X 7 days	
		TRICHOMONIASIS And BACTERIAL VAGINOSIS	□ Metronidazole (Flagyl) 2 gms PO X 1 □ Metronidazole (Flagyl) 1 gm PO now; send 1 gm home with patient to be taken in AM. NO ALCOHOL WITH MED.	Note: If alcohol has been consumed in the last 24 hours, Metronidazole (Flagyl) is to be taken at home after waiting 24 hours from last alcoholic beverage.	
		HEPATITIS B Administer if patient is known to be NOT immune or if immune status is not known.	AGE 11-19: Recombivax: 5 mcg/0.5 ml IM	ADULT OVER AGE 19: ☐ Engerix-B: 20 mcg/ml IM	
		Follow up doses 1-2 months and 4-6 months.	HBIG NO LONGER RECOMMENDED	Contraindicated in severe yeast allergy.	
			AccessPak		
			(contains initial 72 hr supply)		
		HIV - PEP	Combivir 1 tablet PO BID (zidovudine 300mg/lamivudine 150 mg)	Higher Risk: Expanded Regimen (with Viracept) Viracept 5 tablets PO BID (nelfinavir 250 mg)	
		R.N.	M D	Date	